

TALK TO YOUR DOCTOR ABOUT STANDING UP TO BAD CHOLESTEROL

When you have known heart disease, it's important to keep track of your bad cholesterol (LDL-C) levels. This guide can help you have an open and honest conversation with your doctor about taking the next step toward lowering your bad cholesterol.

MAKE THE MOST OF YOUR CONVERSATIONS WITH YOUR DOCTOR

Answer these quick questions and share them with your doctor, so together you can decide the best path forward for you.

1	For people with known heart disease, the recommended cholesterol (LDL-C) level is less than 70 mg/dL. Do you know your number? mg/dL	4	What challenges, if any, have you faced when making changes to your lifestyle? Remembering to take my medication on time, as directed
2	On a scale of 1 to 10 (1 = not worried at all, 10 = very worried) how concerned are you about your bad cholesterol?		Improving my diet and increasing my activity level Limiting unhealthy habits, like smoking and drinking alcohol in excess
3	1 2 3 4 5 6 7 8 9 10 What are the top 3 reasons you want to lower your LDL-C?		Dealing with other health issues, such as diabetes or hypertension Worrying about paying for my medication
	- l worry about having a heart attack, stroke, or other serious health problems*		Other
	l try to be as health conscious as possible I do it for my family (spouse, kids, grandkids),	5	Do you feel like you know what else you can do to lower your bad cholesterol?
	who depend on me to be there for them My doctor recommends it		Yes, I know what additional changes I can make No, I need help to better understand what
	Other not known if LEQVIO can decrease problems related to high lesterol, such as heart attacks or stroke.		else I can do Unsure if I need to be doing more

What is LEQVIO?

LEQVIO (inclisiran) is an injectable prescription medicine used along with diet and other lipid-lowering medicines in adults who need additional lowering of "bad" cholesterol (LDL-C) and have known cardiovascular disease and/or heterozygous familial hypercholesterolemia (HeFH), an inherited condition that causes high levels of LDL-C. It is not known if LEQVIO can decrease problems related to high cholesterol, such as heart attacks or stroke.

IMPORTANT SAFETY INFORMATION

The most common side effects of LEQVIO were: injection site reaction (including pain, redness, and rash), joint pain, urinary tract infection, diarrhea, chest cold, pain in legs or arms and shortness of breath.

Please see LEQVIO full <u>Prescribing Information</u>.



SAY HELLO TO LEQVIO® AND THE POSSIBILITY TO TAKE BAD CHOLESTEROL DOWN

2 doses a year given by your doctor.* Proven to lower bad cholesterol by over 50%.*



FITS INTO TWICE-YEARLY DOCTOR VISITS

LEQVIO is an injection given by your doctor or health care provider, so you don't have to inject yourself.



THE MAJORITY OF SIDE EFFECTS WERE MILD TO MODERATE[‡]

The most common side effect was injection site reaction, which was experienced by 8.2% of people taking LEQVIO vs 1.8% placebo patients.

Side effects leading to discontinuation occurred in 2.5% of people taking LEQVIO vs 1.9% with placebo (a substance that doesn't contain any medication).



LEQVIO WORKS WITH YOUR STATIN AND MAY HELP YOU REACH YOUR GOALS

Which of these are of interest to you?

2 doses a year*

Proven to lower cholesterol by over 50% and keep it low[†]

An injection given by your doctor or health care provider

All of the above



You could pay as little as \$0⁵

Go to **LEQVIO.com/savings-and-support** to learn more about LEQVIO affordability, including for those with Medicare Part B and supplemental insurance, or those with commercial or private insurance.

*2 doses a year after 2 initial doses.

[†]With LEQ^VIO, people in clinical studies on a maximally tolerated statin lowered their cholesterol by 50% and kept it low during each 6-month dosing interval vs placebo (a substance that doesn't contain any medication). Individual results may vary.

*The most common side effects of LEQVIO were: injection site reaction (including pain, redness, and rash), joint pain, urinary tract infection, diarrhea, chest cold, pain in leas or arms and shortness of breath.

[§]Limitations apply. The LEQVIO Co-pay Program may include the Co-pay Card, Payment Card (if applicable), and Rebate. Per-treatment maximums and an annual benefit cap apply. Offer not valid under Medicare, Medicaid, TRICARE, VA, DoD, or any other federal or state program. Novartis reserves the right to rescind, revoke, or amend this Program without notice. See complete <u>Terms & Conditions</u>.

IMPORTANT SAFETY INFORMATION

The most common side effects of LEQVIO were: injection site reaction (including pain, redness, and rash), joint pain, urinary tract infection, diarrhea, chest cold, pain in legs or arms and shortness of breath.

These are not all the possible side effects of LEQVIO. Ask your health care provider for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call <u>1-800-FDA-1088</u>.

Please see LEQVIO full Prescribing Information.

LEQVIO and the LEQVIO logo are registered trademarks of Novartis AG. Licensed from Alnylam Pharmaceuticals, Inc.



Novartis Pharmaceuticals Corporation East Hanover, New Jersey 07936-1080



Get support from the LEQVIO Care Program

Once you've been prescribed LEQVIO, you can get access to a dedicated Care Specialist who offers personalized support, medication reminders, healthy living tips, etc.



Visit <u>careprogram.LEQVIO.com</u> or scan this QR code to learn more.

